

Clinical Policy: Sofosbuvir/Velpatasvir/Voxilaprevir (Vosevi)

Reference Number: GA.PMN.25

Effective Date: 9/17

Last Review Date: 7/2021

Line of Business: Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Sofosbuvir/velpatasvir/voxilaprevir (Vosevi[®]) is a fixed-dose combination oral tablet. Sofosbuvir is a nucleotide analog HCV NS5B polymerase inhibitor, velpatasvir is an NS5A inhibitor, and voxilaprevir is an NS3/4A protease inhibitor.

FDA Approved Indication(s)

Vosevi is indicated for the treatment of adult patients with chronic hepatitis C virus (HCV) infection without cirrhosis or with compensated cirrhosis (Child-Pugh A) who have:

- Genotype 1, 2, 3, 4, 5, or 6 infection and have previously been treated with an HCV regimen containing an NS5A inhibitor*;
- Genotype 1a or 3 infection and have previously been treated with an HCV regimen containing sofosbuvir without an NS5A inhibitor**.
 - Additional benefit of Vosevi over sofosbuvir/velpatasvir was not shown in adults with genotype 1b, 2, 4, 5, or 6 infection previously treated with sofosbuvir without an NS5A inhibitor.

Policy/Criteria

It is the policy of health plans affiliated with Centene Corporation[®] that Vosevi is **medically necessary** when the following criteria are met:

I. Approval Criteria

*** Provider must submit documentation (including office chart notes and lab results) supporting that member has met all approval criteria ***

A. Chronic Hepatitis C Infection (must meet all):

1. Diagnosis of chronic HCV infection as evidenced by detectable HCV RNA (ribonucleic acid) levels by quantitative assay in the last 6 months;
2. Age \geq 18 years;
3. Member meets one of the following (a or b):
 - a. If HCV genotype 1, 2, 3, 4, 5 or 6, member has previously been treated with an HCV regimen containing one of the following NS5A inhibitors: daclatasvir, elbasvir, ledipasvir, ombitasvir, pibrentasvir, or velpatasvir;
 - b. If HCV genotype is 1a or 3, member has previously been treated with an HCV regimen containing sofosbuvir with or without any of the following: peginterferon alfa/ribavirin, ribavirin, HCV NS3/4A protease inhibitor (boceprevir, simeprevir or telaprevir);

CLINICAL POLICY

Sofosbuvir/Velpatasvir/Voxilaprevir

4. Member has received ≥ 8 weeks of the prior direct-acting antiviral agent (DAA) regimen from 2a or 2b above, unless virologic failure was determined prior to 8 weeks of therapy;
5. Member must use Mavyret[®] (as a single agent or in combination as indicated below) if member meets one of the following (a, b or c), unless contraindicated or clinically significant adverse effects are experienced (*see Appendix E*):
 - a. HCV genotype is 1, and member has previously been treated with an HCV regimen containing an NS5A inhibitor without an NS3/4A protease inhibitor (i.e., Daklinza, Epclusa, Harvoni);
 - b. HCV genotype is 1a or 3, and member has previously been treated with an HCV regimen containing sofosbuvir with or without any of the following: peginterferon alfa/ribavirin, ribavirin, HCV NS3/4A protease inhibitor (boceprevir, simeprevir or telaprevir);
 - c. For HCV genotype 1 through 6 and member previously treated with Vosevi, Mavyret must be used in combination with Sovaldi[®] and RBV;
6. Life expectancy ≥ 12 months with HCV treatment;
7. Prescribed regimen is consistent with an FDA or AASLD-IDSAs recommended regimen (*see Appendix D Dosage*);
8. Member is hepatitis B virus (HBV) negative, or if positive, documentation that concurrent HBV infection is being treated (e.g., tenofovir alafenamide, adefovir, entecavir), unless contraindicated or clinically significant adverse effects are experienced (*see Appendix E*);
9. Prescribed dose does not exceed one tablet (sofosbuvir 400 mg/velpatasvir 100 mg/voxilaprevir 100 mg) daily.

Approval duration: up to a total of 24 weeks*

(*Approved duration should be consistent with a regimen in Appendix D FDA approved dosages and Treatment Duration)

B. Other diagnoses/indications: Refer to CP.PMN.53 – No Coverage Criteria/Off-Label Use Policy if diagnosis is NOT specifically listed under section I.

II. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

AASLD: American Association for the Study of Liver Diseases	MRE: magnetic resonance elastography
APRI: AST to platelet ratio	NS3/4A, NS5A/B: nonstructural protein
CTP: Child Turcotte Pugh	Peg-IFN: pegylated interferon
CrCl: creatinine clearance	PI: protease inhibitor
FDA: Food and Drug Administration	RBV: ribavirin
FIB-4: Fibrosis-4 index	RNA: ribonucleic acid
HCC: hepatocellular carcinoma	
HCV: hepatitis C virus	
IDSAs: Infectious Diseases Society of America	

CLINICAL POLICY

Sofosbuvir/Velpatasvir/Voxilaprevir

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
Mavyret® (glecaprevir/ pibrentasvir)	Treatment-experienced with IFN/pegIFN, RBV and/or sofosbuvir: Genotypes 1, 2, 4, 5, or 6 Without cirrhosis: Three tablets PO QD for 8 weeks With compensated cirrhosis: Three tablets PO QD for 12 weeks	Mavyret: glecaprevir 300 mg/ pibrentasvir 120 mg (3 tablets) per day
Mavyret® (glecaprevir /pibrentasvir)	Treatment-experienced with IFN/pegIFN, RBV and/or sofosbuvir: Genotype 3 Without cirrhosis or with compensated cirrhosis: Three tablets PO QD for 16 weeks	Mavyret: glecaprevir 300 mg/ pibrentasvir 120 mg (3 tablets) per day
Mavyret® (glecaprevir /pibrentasvir)	Treatment-experienced with NS5A inhibitor without prior NS3/4A protease inhibitor: Genotype 1 Without cirrhosis or with compensated cirrhosis: Three tablets PO QD for 16 weeks	Mavyret: glecaprevir 300 mg/ pibrentasvir 120 mg (3 tablets) per day
Mavyret® (glecaprevir /pibrentasvir)	Treatment-experienced with NS3/4A protease inhibitor without prior NS5A inhibitor: Genotype 1 Without cirrhosis or with compensated cirrhosis: Three tablets PO QD for 12 weeks	Mavyret: glecaprevir 300 mg/ pibrentasvir 120 mg (3 tablets) per day
Mavyret®(glecaprevir /pibrentasvir) + Sovaldi® (sofosbuvir) + RBV	With prior sofosbuvir/velpatasvir/voxilaprevir treatment failure, with compensated cirrhosis or without cirrhosis” Genotypes 1-6 †: Sovaldi 400 mg + Mavyret 300 mg/120 mg + weight-based RBV for 16 weeks	Varies

Therapeutic alternatives are listed as Brand Name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

† Off-label, AASLD-IDSA guideline-supported dosing regimen

CLINICAL POLICY

Sofosbuvir/Velpatasvir/Voxilaprevir

Appendix C: Contraindications

- Coadministration with rifampin
- Boxed warning(s): risk of hepatitis B virus reactivation in patients coinfecting with HCV and HBV

Appendix D: Direct-Acting Antivirals for Treatment of HCV Infection

Brand Name	Drug Class				
	NS5A Inhibitor	Nucleotide Analog NS5B Polymerase Inhibitor	Non-Nucleoside NS5B Palm Polymerase Inhibitor	NS3/4A Protease Inhibitor (PI)**	CYP3A Inhibitor
Daklinza	Daclatasvir				
Epclusa*	Velpatasvir	Sofosbuvir			
Harvoni*	Ledipasvir	Sofosbuvir			
Mavyret*	Pibrentasvir			Glecaprevir	
Sovaldi		Sofosbuvir			
Viekira /PAK*	Ombitasvir		Dasabuvir	Paritaprevir	Ritonavir
Vosevi*	Velpatasvir	Sofosbuvir		Voxilaprevir	
Zepatier*	Elbasvir			Grazoprevir	

*Combination drugs

Appendix E: General Information

- Hepatitis B Virus (HBV) Reactivation is a black box warning for all direct-acting antiviral drugs for the treatment of HCV. HBV reactivation has been reported when treating HCV for patients co-infected with HBV, leading to fulminant hepatitis, hepatic failure, and death, in some cases. Patients should be monitored for HBV reactivation and hepatitis flare during HCV treatment and post-treatment follow-up, with treatment of HBV infection as clinically indicated.
 -
- Unacceptable medical justification for inability to use Mavyret (preferred product):
 - Black Box warning (BBW): currently or previously infected with hepatitis B virus. This BBW is not unique to Mavyret, and it applies across the entire therapeutic class of direct-acting antivirals for treatment of HCV infection. Therefore it is not a valid clinical reason not to use Mavyret.
 - Concurrent anticoagulant therapy: Fluctuations in International Normalized Ratio (INR) have been observed in warfarin recipients who were also receiving treatment for HCV infections. This BBW is not unique to Mavyret, and it applies across the entire therapeutic class of direct-acting antivirals for treatment of HCV infection..
 - Drug-drug interactions with one or more of the following agents:
 - Rifampin, carbamazepine, or St. John's wort:

CLINICAL POLICY

Sofosbuvir/Velpatasvir/Voxilaprevir

- These drug-drug interactions are not unique to Mavyret, and they apply across the entire therapeutic class of direct-acting antivirals for the treatment of HCV infection.

III. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose	Reference
Genotype 1-6: Treatment-experienced with NS5A inhibitor* with or without compensated cirrhosis	One tablet PO QD for 12 weeks	One tablet (sofosbuvir 400 mg/ velpatasvir 100 mg/ voxilaprevir 100 mg) per day	1) FDA-approved labeling 2) AASLD-IDSA (updated November 2019)
Genotype 1a or 3: Treatment-experienced with a sofosbuvir- containing regimen without NS5A inhibitor* with or without compensated cirrhosis	One tablet PO QD for 12 weeks		1) FDA-approved labeling 2) AASLD-IDSA (updated November 2019)
Genotype 1-6: Treatment-experienced with Vosevi® with or without compensated cirrhosis	Vosevi one tablet PO QD with weight- based RBV for 24 weeks		AASLD-IDSA (updated November 2019)

AASLD/IDSA treatment guidelines for chronic hepatitis C infection are updated at irregular intervals; refer to the most updated AASLD/IDSA guideline for most accurate treatment regimen.

* See appendix E

IV. Product Availability

Tablet: sofosbuvir 400 mg/velpatasvir 100 mg/voxilaprevir 100 mg

V. References

1. Vosevi Prescribing Information. Foster City, CA: Gilead Sciences, Inc.; November 2019. Available at: www.vosevi.com. Accessed April 15, 2021.
2. American Association for the Study of Liver Diseases/ Infectious Disease Society of America (AASLD-IDSA). HCV guidance: recommendations for testing, managing, and

CLINICAL POLICY

Sofosbuvir/Velpatasvir/Voxilaprevir

treating hepatitis C. Last updated March 12, 2021. Available at: <https://www.hcvguidelines.org/>. Accessed April 15, 2021.

3. Bourliere M, et al. Sofosbuvir, velpatasvir, and voxilaprevir for previously treated HCV infection. NEJM 2017;376:2134-46.
4. CDC. Hepatitis C Q&As for health professionals. Last updated August 7, 2020. Available at: <https://www.cdc.gov/hepatitis/hcv/hcvfaq.htm>. Accessed April 15, 2021.

Reviews, Revisions, and Approvals	Date	Approval Date
Policy created	09/17	9/17
Changed current Georgia policy templates to corporate standard templates for drug coverage criteria to meet corporate compliance. Changes/revisions included; new formatting, font size, use of standard policy language for each section of policy, and rearranged order of certain steps in criteria and sections. Added new preferred treatment tables that includes dosage and frequency based on genotype for Mavyret. Removed background sections. Updated general information and contraindication section to be consistent with corporate HCV policies.	2/21/19	2/19
Annual review. Updated wording of approval criteria for meeting use of preferential Mavyret.	10/19	10/19
Added Epclusa to Appendix B. Removed Appendix D for Metavir scoring. Removed dosing for Mavyret treatment-naïve. Updated order of all other Appendices. Updated references.	4/2020	4/2020
Added preferred re-direction for off-label Mavyret + Sovaldi + RBV after Vosevi failure; modified initial and continued approval durations up to 24 weeks to allow for post Vosevi failure off-label indication dosing per per AASLD/IDSA guideline; references reviewed and updated.	7/2020	7/2020
Annual review. Added “quantity assay” specifically for testing to initial criteria. Added that preferred Mavyret needs to be used as a single agent or in combination as indicated to initial criteria. Updated criteria to include pibrentasvir as an acceptable option for previous treatment with an HCV regimen containing an NS5A inhibitor to align with appendix D table; references reviewed and updated. Added additional Mavyret plus RBV dosing regimen to Therapeutic Alternatives table. Removed “chronic HCV infection” verbiage from Therapeutic Alternative table to be consistent with CNC policy. Added Box Warning for hepatitis B reactivation to Appendix C: Contraindications. Added Mavyret and Vosevi to Appendix D-Direct Acting Antivirals for Treatment of HCV infection and removed Olysio, Technivie, and Viekira XR as these were previously removed from the market. Changed Centene Logo to PSHP Logo. Minor formatting changes throughout document. References reviewed.	4/2021	4/2021
Updated Appendix B therapeutic alternatives; removed the appendix E acceptable medical justification section for inability to use Mavyret as it	7/2021	7/2021

CLINICAL POLICY

Sofosbuvir/Velpatasvir/Voxilaprevir

Reviews, Revisions, and Approvals	Date	Approval Date
overlaps with Vosevi clinical parameters for not using; references reviewed and updated.		

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

CLINICAL POLICY

Sofosbuvir/Velpatasvir/Voxilaprevir

Note: For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.